

Color quality improvement of Cherax quadricarinatus with dragon fruit peel meal utilization

by Adi Candra

Submission date: 03-Jun-2024 02:26PM (UTC+0700)

Submission ID: 2295146316

File name: Rakhmawati_et_al_Des_2023_Jurnal_Depik.pdf (344.36K)

Word count: 4355

Character count: 22277



Color quality improvement of *Cberax quadricarinatus* with dragon fruit peel meal utilization

Rakhmawati Rakhmawati^{1*}, Tulas Aprilia², Nur Indariyanti³, Andre Saputra⁴, Eulis Marlina⁵

¹Department of Aquaculture, Lampung State Polytechnic, Lampung, Indonesia

²Department of Fish Hatchery Technology, Lampung State Polytechnic, Lampung, Indonesia

³Department of Fish Hatchery Technology, Lampung State Polytechnic, Lampung, Indonesia

⁴Department of Fish Hatchery Technology, Lampung State Polytechnic, Lampung, Indonesia

ARTICLE INFO

Keywords:

Dragon fruit peel meal
 Utilization
 Diet
 Color quality
Cberax quadricarinatus

DOI: 10.13170/depik.12.3.28628

10

ABSTRACT

This study aims to determine the effect of dragon fruit peel meal utilization in the diet to improving the color quality of freshwater crayfish (*Cberax quadricarinatus*). This research was done from May to June 2022 at the Lampung State Polytechnic Fisheries Laboratory. The test animals used were freshwater crayfish with an average weight of 6.7 – 6.8 g/crayfish and a stocking density of 7 crayfish/aquarium. Feeding with a feeding rate of 3% twice a day. The experimental design used was a completely randomized design with 5 treatments and 3 repetitions. The treatments were A = commercial feed (control), B = 5% dose, C = 10% dose, D = 15% dose, E = 20% dose. The parameters observed in this study were an increase in color quality and water quality. The results of this study indicate that the difference in the dose of red dragon fruit peel meal in the diets can have a significant effect on the color brightness of freshwater crayfish (7.01 in claws and 87.59 in Cephalotorax to teston). The conclusion of this research is that supplementation of red dragon fruit peel meal mainly at a dose of 10% in the feed can improve the color quality of freshwater crayfish from brown to bluish color.

Introduction

The high demand for red clawed freshwater crayfish as an ornamental aquaculture commodity and as a consumption commodity in urban areas is a great opportunity to increase the productivity of its cultivation. Meanwhile, the availability of red clawed freshwater crayfish is still limited. Thus the price is still relatively high and is not available all the time. Other side, during the COVID-19 pandemic in recent years, the consumption of dragon fruit by the Indonesian people has increased, including in Lampung Province (BPS, 2019). Along with public awareness of the importance of increasing the vitality of the body by consuming fruit. This increase in consumption also causes the waste of dragon fruit peel to the environment to increase. This needs to be handled so that it can be utilized and the environment can be kept clean due to the reduction of the waste.

Dragon fruit peel waste contains essential nutrients needed by fish or shrimp or crayfish to increase the productivity of their culture, especially through diets composition engineering technology. This can be a solution for feed producers and increase profits for freshwater crayfish (*Cberax quadricarinatus*) farmers.

One of the raw materials that can improve color quality and is a food waste is dragon fruit peel. Dragon fruit contains betacyanin, which is a red-violet pigment (Faridah *et al.*, 2014), and also contains anrospanin, which is a red to orange pigment (Simanjuntak *et al.*, 2014; Ramadhani *et al.*, 2017). In addition, dragon fruit peel contains antioxidants, namely flavonoids, phenolics, and carotenoids (Wahdaningsih *et al.*, 2017; Pujiastuti and ElZeba, 2021). These antioxidants are able to inhibit the activity of free radicals (Akbar *et al.*, 2023). Rakhmawati (2017) reported that dragon fruit peel

* Corresponding author.

Email address: rakhmawati@polinela.ac.id

inhibits free radical activity by 50% by 397.64 ppm and contains vitamin C by 9.79 ppm. Aprilia and Rakhmawati (2021) reported that in terms of nutrients, fruit peel flour contains 5.08% protein; carbohydrates 50.74%; crude fiber 26.22%; ash 21.35% and fat 4.80% (wet weight).

The challenge of red claw crayfish rearing so far is to improve the color quality when developed on certain media. This needs to be solved, considering that aquaculture commodities cannot synthesize color pigments in their bodies, so they must be fed from outside. In addition, low survival and slow growth are important issues to find a solution to increase the availability of this commodity and efforts are needed to improve the color quality. The use of dragon fruit peel meal has never been studied in red clawed freshwater crayfish (*C. quadricarinatus*). This can be achieved by using certain nutrients that support the color quality and health of red clawed freshwater crayfish. The results of research on the nutrient content of dragon fruit peel meal and its positive effect on improving the quality of fish color so it is necessary to do applied research in the application of freshwater crayfish culture.

⁸
Materials and Methods

Preparation of dragon fruit peel meal

Dragon fruit peel meal was prepared according Aprilia and Rakhmawati (2021). The Preparation of dragon fruit peel meal begins with cleaning and washing the skin and the scales. The dragon fruit peel was weighed and cut into small pieces, then dried in an oven at 23°C for 7 hours. After drying, it was milled with a blender and sieved using a 60 mesh sieve to obtain red dragon fruit peel meal. Dried dragon fruit peel meal will be mixed with other raw materials to make the experimental feed.

Preparation the experimental diets ¹

The feed to be made contains 38% protein. The ingredients and composition of the test feed are presented in Table 1. This study used five types of test feed containing dragon fruit peel flour with different doses. The test feed was substituted with a dose of 0 (control, without dragon fruit peel flour); 5%; 10%; 15% and 20%.

All raw materials are weighed and mixed evenly, followed by the addition of oil and water. The feed was molded with a diameter of 1 mm, dried in a tumble dryer, and stored in plastic containers until use.

Experimental animal and rearing

Two hundred red claw freshwater crayfish juvenile with an initial length of 5-6 cm and an initial weight of 6.7 – 6.8 g were obtained from Frandy

²⁸
Farm Freshwater Lobster, Natar District, South Lampung Regency, Lampung Province, Indonesia.

Table 1. Composition of feed ingredients with dragon fruit peel meal (DFPM) utilization

Ingredient (g kg ⁻¹)	Supplementation of dragon fruit peel meal in the diets (g kg ⁻¹)				
	0	5%	10%	15%	20%
Fish meal	180	180	180	180	180
Soybean meal	350	350	350	350	350
DFPM	0	50	100	150	200
Cornstarch	300	250	200	150	100
Fish oil	50	50	50	50	50
Corn oil	50	50	50	50	50
CMC	20	20	20	20	20
Vitamin & Mineral	50	50	50	50	50
	1000	1000	1000	1000	1000

Prior to the study, red clawed crayfish were kept for 1 week and given commercial feed (35% protein) for acclimatization to the research conditions. Previously, the shrimp were adapted to the rearing medium for approximately 3 weeks to uniform size. The test red claw lobster were reared for 28 days. At the beginning of the study, individual lobsters were weighed and randomly distributed into 15 aquariums (40 x 40 x 40 cm³) with a density of 7 lobsters per aquarium. Feeding with a feeding rate of 3% twice a day. Continuous aeration was provided with 25% water change every 24 hours. Red claw lobster feces are siphoned from the tub at 16.00 every day. Water quality parameters in all experimental media include temperature, dissolved oxygen, total ammonia nitrogen and pH. Evaluate the lobster survival rate every week of the rearing period. The survival rate was calculated by the formula Survival Rate = (Number of test lobsters at the end of rearing (tails) / Number of test lobsters at the beginning of rearing (tails)) x 100%.

Evaluation of color change of red claw lobster

Observation of lobster body color was carried out once every 14 days for 4 weeks Parameters observed in the study include the main parameters. The main parameter is the observation of red claw body color every 14 days using the scoring method by determining the red claw body color scale based on the Trumatch color standard, as applied in the study Amin et al., (2012). The color scale used is scale 1: light brown, scale 2: medium brown, scale 3: dark brown, scale 4: light green, scale 5: medium green,

scale 6: dark green, scale 7: light blue, scale 8: medium blue, scale 9: dark blue.

The observations were carried out by three people to prevent deviations. Parameters observed were changes in the color of crayfish. The increase in color was observed by comparing the colors of the two parts, namely the claws and cephalothorax to the telson, with a color standard.




Figure 1. Trumatch color standard

Data analysis

The study was conducted with five treatments and three replications, namely treatment A without dragon fruit peel powder treatment (control), treatment B, namely dragon fruit peel flour treatment dose of 5%, treatment C, dragon fruit peel flour treatment dose of 10%, treatment D, dose treatment 15% dragon fruit peel flour, 20 treatment E, namely the dose treatment of 20% dragon fruit peel flour. The data obtained in this study were statistically analyzed using the one way ANOVA test and the BNJ (Tukey) follow-up test ($p < 0.05$).

Results

Color Changes in Freshwater Lobster Red claw (*Cherax quadricarinatus*). Color observations included two parameters, namely color observations on the claws and cephalothorax to the freshwater crayfish telson (*Cherax quadricarinatus*). During the observation, an increase in color quality began to appear on the 14th day. Based on observations, the highest average value was 83.61 in the claws and 85.55 in the Cephalothorax section until telson began to appear in treatment C (10% dose) after 14 days maintenance. Color scoring can be seen in Table 2.

The results of observations for 28 days, the color of the red claw crayfish changes every week, except for the first week because the crayfish are still adapting to the feed given. The increase in color quality every week is caused by the provision of a color pigment source, this indicates that there is a difference in color quality improvement in freshwater crayfish in the aquarium. The results showed that there was an increase in the quality of lobster color quality in treatment C (10% Red

Dragon Fruit Peel Meal) with an average sample value of 86.01 for claws and 87.9 for chepalotorax to teslon.

Table 2. Scoring of increased brightness of claws and cephalothorax to telson red claw (*Cherax quadricarinatus*) for 3 observations in 28 days of rearing.

Observed part	Treatment	Average Observation Score (Day-)		
		0	14	28
Claw	A	79.36	80.55	80.66
	B	77.77	78.95	83.69
	C	76.19	83.61	86.01
	D	77.77	80.13	83.69
	E	77.77	80	82.85
Cephalothorax up telson	A	74.60	80.55	84.25
	B	73.01	81.66	82.87
	C	73.01	85.55	87.59
	D	73.01	85.55	86.50
	E	74.60	82.03	83.33

The increase in color quality in treatment D was not significantly different from treatment C, which ranged from 83.69 to claws and 86.50 to telson. The score in treatment B (Dose of 5%) was 83.69 on the skin and 82.87 on the Cephalothorax section, then the score on treatment E (20% dose) was 82.85 on the claws and 83.33 on the Cephalothorax up to telephone. While the lowest score is in treatment A (Control) with a value of 80.66 in the claw section and 84.25 in the Cephalothorax section to telson.

Table 3. Water quality during rearing

Treatments	Dissolve Oxygen (mg l^{-3})	Temperature ($^{\circ}\text{C}$)	pH	Total ammonia nitrogen (mg l^{-3})
A	6.4	27.4	8.23	0.25
B	6.6	27.6	8.21	0.25
C	6.9	27.2	8.11	0.15
D	6.7	27.5	8.17	0.15
E	6.4	27.4	8.21	0.25

The survival rate of freshwater crayfish for 28 days of rearing is in treatment A (control) of 90.47%, treatment B with a dose of 5% at 85.71%, treatment C with a dose of 10% at 71.42%, treatment D with a dose of 15% of 80.95% and treatment E of 80.95%. Water quality includes dissolved oxygen (DO), water

temperature, acidity (pH), and ammonia could be seen in Table 3.

Discussion

The results of observations for 28 days, the color of the red claw crayfish changes every week, except for the first week because the crayfish are still adapting to the feed given. The increase in color quality every week is caused by the provision of a color pigment source, this indicates that there is a difference in color quality improvement in freshwater crayfish in the aquarium. Changes in the color of the crayfish began to occur on the 7th and 14th day with the addition of 5% red dragon fruit peel flour to the feed due to an increase in carotenoids in pigment cells (Simamora, 2019). Meanwhile, in sand lobsters, their entire body color changed after 97 days of rearing when they were fed spirulina-added feed (Soffa et al., 2019).

In addition, the results showed that there was an increase in the quality of red claw crayfish color quality in treatment C (10% red dragon fruit peel meal) with an average sample value of 86.01 for claws and 87.9 for chepalotorax to teslon. The increase in color quality in treatment D was not significantly different from treatment C, which ranged from 83.69 to claws and 86.50 to telson. The score in treatment B (Dose of 5%) was 83.69 on the skin and 82.87 on the Chepalotorax section, then the score on treatment E (20% dose) was 82.85 on the claws and 83.33 on the Chepalotorax up to telephone. While the lowest score is in treatment A (Control) with a value of 80.66 in the claw section and 84.25 in the chepalotorax section to teslon.

The color appears first from the tip of the claw which then spreads to the base of the claw. Improving the quality of red claw body color is influenced by the amount of carotenoids contained in the exoskeleton. Beta-carotenoid, vitamins and anthocyanin in dragon fruit peel is thought to increase the skin color lightening of crayfish. The beta-carotenoid contained in it fulfills the needs of the crayfish body, because it cannot synthesize itself (Nururrahmah and Widiarnu, 2013; Kalidupa et al., 2018). Kembuan et al. (2012) and Noor et al. (2016) stated that the vitamin C contained in it acts as a strong antioxidant and is absorbed easily in the skin. The antioxidant content in super dragon fruit peel extract reaches 30 b/v (Ni'matusyukriyah and Sarwono, 2019) and even reaches 58 mg/l (Putri et al., 2015). Anthocyanins act as nutrients that can provide an orange color and color strengthening (Priska et al., 2018). The phenolic activity contained as an antioxidant is also thought to play a role in this.

Lorith and Kanlayavattanakul (2013) stated that phenolic activity is the basis of an antioxidant mechanism that stabilizes pH and temperature and is responsible for brightening colors (Pramitasari et al., 2022). The contents of beta-carotenoid, vitamins and antioxidants in dragon fruit peel meal, so the color will also first appear on that part of the body. After the color appears at the tips of the claws and cephalothorax, then the color spreads throughout the claws and abdomen to the telson.

Several previous research results coloring ornamental fish using natural dyes with the addition of red dragon fruit peel meal in feed can increase the brightness of color in ornamental fish. This is proven by research conducted by Kalidupa et al. (2018) the administration of red dragon fruit peel meal in the feed significantly affected the increase in the brightness of the orange and black colors in koi carp. In addition, giving red dragon fruit peel flour with a dose of 15% in feed can increase the brightness of the orange and black colors in koi carp (Kalidupa et al., 2018). Color improvement in koi fish fed 10% dragon fruit meal (Kurnia et al., 2019) and in red cherry shrimp added astaxanthin and canthaxanthin (Amin et al., 2012). Apart from affecting the color of the dragon fruit skin, it can also affect health as a natural antibiotic that can suppress the attack of *Aeromonas hydrophilla* bacteria on tilapia (Andayani et al., 2018; Sudrajat, 2019) and increase growth in goldfish (Efianda et al., 2018) and in snakehead fish (Rakhmawati et al., 2021).

This study was conducted using raw materials in the manufacture of feed by mixing red dragon fruit peel flour as natural dye. Each treatment had a different dose of red dragon fruit peel flour so that the proximate content was different. Based on the results of observations showed that each treatment there was an increase in the color quality of the lobster. The highest increase in color quality was found in lobsters fed C feed (10% dose). Followed by treatment D (15% dose), treatment E (20% dose), treatment B (5% dose) and the lowest was treatment A (without red dragon fruit skin flour). excessive and not lacking to obtain an increase in color quality in red claw lobster. Amiin et al. (2012) argued that to obtain optimal color improvement in red cherry shrimp by adding 100 mg/kg canthaxanthin. The same thing with rainbow trout, Meilisza et al. (2021), Kurumoi rainbowfish fed 2.6 g/kg asthaxanthin supplementation had higher enhance color than 5.1 g/kg.

Color quality improvement in red claw crayfish (*C. quadricarinatus*) is influenced by several internal and external factors. Internal factors include species,

gender, organs and hormones. But for this study we did not select the sex at the start of rearing considering the small size of the red claw lobsters. External factors include background color of the rearing place, light intensity, temperature of rearing conditions and carotenoids in the feed. Other external factors besides carotenoids in the feed in the form of background color of the rearing place, light intensity and temperature of rearing conditions that affect the increase in red claw body color are important factors that also need to be considered (Diaz – Jimenez et al., 2018).

Water quality parameters are still within the optimal range of rearing. So is the survival rate during maintenance. The survival rate can be said to be still high. Samad et al. (2022) stated that the survival rate of freshwater crayfish ranged from 80 to 93.33% which was still high. However, treatment C was still less than optimal due to cannibalism in freshwater crayfish during maintenance. The high survival rate of freshwater crayfish is due to both the amount and timing of feeding as well as the condition of water quality during maintenance which does not become a limiting factor in the survival of freshwater crayfish.

Conclusion

The results showed that dragon fruit peel meal improve the color of freshwater crayfish (*Cherax quadricarinatus*). The brightest color was lobster which was given a utilization of 10% dragon fruit peel meal from brown to a bluish color.

Acknowledgments

Thank you to Lampung State Polytechnic which has provided funds for this research through DIPA for Fiscal Year 2022 with Number: 321.59/PL15.8/PP/2022 on the Applied Research Scheme.

References

Akbar, S.A., M. Hasan, S. Afriani, C. Nuzlia. 2023. Evaluation of phytochemical composition and metabolite profiling of macroalgae *Caulerpa taxifolia* and *C. peltata* from the Banda Aceh coast, Indonesia. *Biodiversitas Journal of Biological Diversity*, 24(10): 5283-5292.

Amin M.I., Rosidah, W. Lili. 2012. Peningkatan kecerahan warna udang red cherry (*Neocaridina heteropoda*) jantan melalui pemberian astaxanthin dan canthaxanthin dalam pakan. *Jurnal Perikanan dan Kelautan*, 3(4): 243 – 252.

Andayani, S., H. Suprastyani, I. Masfiah. 2018. Pengaruh pemberian ekstrak kasar kulit buah naga (*Hylocereus costaricensis*) terhadap histologi hati ikan nila yang terinfeksi *Aeromonas hydrophilla*. *Journal of Fisheries and Marine Research*, 3(2): 149 – 159.

Aprilia T, R. Rakhmawati. 2021. Peningkatan kualitas komposisi kimia pakan dengan penambahan tepung kulit buah naga (*Hylocereus polyrhizus*). *Jurnal Rekayasa dan Teknologi Budidaya Perairan*, 9(2): 1101-1108.

BPS. Badan Pusat Statistik. Statistik tanaman buah-buahan dan sayuran-tahunan.

<http://www.bps.go.id/publication/2019/10/07/1846605363955649e8f6dd6d/>. 2019.

Diaz-Jimenez, J., M.P. Hernandez-Vergara, C.I. Perez-Rostro. 2018. The effect of background colour and lighting of aquarium on the body pigmentation of peppered shrimp *Lyssmata wurdemanni*. *Aquaculture Research*, 49: 3508 – 3516.

Efianda T.R., Y. Yusnita, N. Najmi, K.R. Ananda, F. Saputra. 2020. Pengaruh kulit buah naga (*Hylocereus polyrhizus*) dalam pakan terhadap kinerja reproduksi ikan koi (*Cyprinus carpio*). *Jurnal Perikanan Tropis*, 7(2): 107-113.

Kalidupa N, A. Kurnia, I. Nur. 2018. Studi pemanfaatan tepung kulit buah naga merah (*Hylocereus polyrhizus*) dalam pakan terhadap pewarnaan ikan koi (*Cyprinus carpio* L.). *Media Akuatika*, 3(1): 590-597.

Kembuan, M.V., S. Wangko, G.N. Tanudjaja. 2012. Peran vitamin C terhadap pigmentasi kulit. *Jurnal Biomedik*, 4 (3): 13 – 17.

Kurnia, A., I. Nur, W.H. Muskita, M. Hamzah, W. Iba, R.S. Patadjai, A.M. Balubi, N. Kalidupa. 2019. Improving skin coloration of koi carp (*Cyprinus carpio*) fed with red dragon fruit peel meal. *AAAL Bioflux*, 12 (4): 1045 – 1053.

Lourith, N. M. Kanlayattanakul. 2013. Antioxidant and stability of dragon fruit peel colour. *Agro food Industry High Technology*, 24(3): 55 – 58.

Meilisza, N., M.A. Suprayudi, D. Jusadi, M.Zairin, I.M. Artika. 2021. Effects of synthetic astaxanthin, chlorella, and spirulina supplementation in diets on growth and pigmentation kurumi rainbowfish, *Melanotaenia parva*. *Indonesian Aquaculture Journal*, 15(2): 67 – 75.

Ni'matusyukriyah, M.A.H. Swasono. 2019. Pengaruh fortifikasi ekstrak kulit buah naga super merah (*Hylocereus costaricensis*) terhadap kandungan antioksidan tape singkong kuning (*Manihot utilisima* Pohl). *Teknologi Pangan*, 11(1):52-65.

Noor M.I., E. Yufita, Zulfalina. 2016. Identifikasi kandungan ekstrak kulit buah naga merah menggunakan Fourier Transform Infrared (FTIR) dan fitokimia identification content of the red dragon fruit extract skin using Fourier Transform Infrared (FTIR) and phytochemistry. *Journal of Aceh Physics Society (JAcPS)*, 5(1): 14–16.

Kalidupa N, A. Kurnia, I. Nur. 2018. Studi pemanfaatan tepung kulit buah naga merah (*Hylocereus polyrhizus*) dalam pakan terhadap pewarnaan ikan koi (*Cyprinus carpio* L.). *Media Akuatika*, 3(1): 590-597.

Nururrahmah, W. Widiarnu. 2013. Analisis kadar beta-karoten kulit buah naga menggunakan spektrofotometrik. *Jurnal Dinamika*, 4 (1): 15 – 26.

Pramitasari, L.N. Gunawicahya, D.S.B Anugrah. 2022. Development of an indicator film based on cassava starch–chitosan incorporated with red dragon fruit peel anthocyanin extract. *Polymers*, 14 (19): 4142.

Priska, M., N. Peni, L. Carvallo, Y.D. Ngapa. 2018. Review: antosianin dan pemanfaatannya. *Cakra Kimia*, 6(2): 79.

Pujastuti, E., D. El'Zeba. 2021. Perbandingan kadar flavonoid total etanol 70% dan 96% kulit buah naga merah (*Hylocereus polyrhizus*) dengan spektrofotometri. *Cendia Journal of Pharmacy*, 5(1): 28 – 43.

Putri, N.K.M., I.W.G. Gunawan, I.W. Suarsa. 2015. Aktivitas antioksidan antosianin dalam ekstrak etanol kulit buah naga super merah (*Hylocereus costaricensis*) dan analisis kadar total. *Jurnal Kimia*, 9 (2): 243 – 251.

Rakhmawati, M. 2017. Aktivitas antioksi dan ekstrak kulit buah naga merah (*Hylocereus polyrhizus*) secara in vitro. S Si. thesis, Universitas Jember, Jember.

Rakhmawati, R., T. Aprilia, A. Kurniawan. 2021. Enhancement the growth of snakehead (*Channa striata*) with addition of dragon fruit peel flour to the diet. *Sriwijaya Journal Environment*. 6 (2): 53-58.

Ramadhani, N.S. 2017. Mikroenkapsulasi ekstrak betasianin dari kulit buah naga merah (*Hylocereus polyrhizus*) menggunakan bahan penyalut maltodekstrin yang dikombinasikan dengan gum arab, cmc, dan karegenan. Magister thesis, Universitas Brawijaya.

Samad, A.P.A., M.F. Isma, E. Ayuzar, Ilhamdi, R. Humairani, F. Santi. 2022. Growth and survival rate of freshwater lobster (*Cherax*

Rakhmawati et al. (2023)

- quadricarinatus*) fed different forage feed. IOP Conference Series: Earth Environment Science 956 012006
- Simamora, D. Pengaruh konsentrasi tepung kulit buah naga merah (*Hylocereus polyrhizus*) pada pakan terhadap peningkatan warna dan pertumbuhan ikan badut (*Ampiphrion ocellaris*). Undergraduate (S1) thesis, University of Borneo Tarakan. 2019.
- Simanjuntak, L., C. Sinaga, Fatimah. 2014. Ekstraksi pigmen antosianin dari kulit buah naga merah (*Hylocereus polyrhizus*). Jurnal Teknik Kimia USU, 3 (2):25.
- Soffa, F.B., I.S. Pratama, A. Ridwanudin, V. Fahmi. 2019. Color alteration and growth performance of spiny lobster (*Panulirus homarus*) juveniles fed with different spirulina concentration in formulated diet. Oseanologi dan Limnologi di Indonesia, 4(2):101-112.
- Sudrajat N. Pengaruh pemberian ekstrak kulit buah naga (*Hylocereus polyrhizus*) terhadap perubahan histopatologi ginjal ikan nila (*Oreochromis niloticus*) yang diinfeksi bakteri *Aeromonas hydrophila*. Undergraduate (S1) thesis, University of Muhammadiyah Malang. 2019.
- Wahdaningsih, S., S. Wahyuono, S. Riyanto, R. Murwanti. 2017. Penetapan kadar fenolik total dan flavonoid total ekstrak methanol dan fraksi etil asetat kulit buah naga merah (*Hylocereus polyrhizus* (F.A.C.WEBER) BRITTON DAN ROSE), Jurnal Ilmiah Farmasi, 6 (3): 295 – 301.

How to cite this paper:

Rakhmawati, R., T. Aprilia, N. Indariyanti, A. Saputra, E. Marlina. 2022. Color Quality Improvement of *Cherax quadricarinatus* with Dragon Fruit Peel Meal Utilization. Depik Jurnal Ilmu-Ilmu Perairan, Pesisir dan Perikanan, 12(3): 314-319.

Color quality improvement of Cherax quadricarinatus with dragon fruit peel meal utilization

ORIGINALITY REPORT

16%

SIMILARITY INDEX

12%

INTERNET SOURCES

11%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

- 1** R Rakhmawati, E Marlina, W Warji. " Efficacy Dietary Supplementation of Banana Peel Meal on Growth and Cannibalism level of Giant Freshwater Prawn ", IOP Conference Series: Earth and Environmental Science, 2020
Publication 2%
- 2** Submitted to unikal
Student Paper 2%
- 3** Submitted to Syiah Kuala University
Student Paper 1%
- 4** R A Prastiya, AM Yusuf. "Red dragon fruit peel (Hylocereus polyrhizus) as feed biosupplement on performance of fattening rabbits", IOP Conference Series: Earth and Environmental Science, 2021
Publication 1%
- 5** repository.umi.ac.id
Internet Source 1%
- 6** jurnal.unsyiah.ac.id
Internet Source 1%

7	journal.unhas.ac.id Internet Source	1 %
8	jurnal.unpad.ac.id Internet Source	1 %
9	JP Batubara, K Laila, R Rumodang, D Kurniawan. "The Effects of Different Dosages of Pasak Bumi (Eurycoma longifolia Jack) towards The Sex of Betta Fish Embryo", IOP Conference Series: Earth and Environmental Science, 2022 Publication	1 %
10	Submitted to Udayana University Student Paper	<1 %
11	www.scilit.net Internet Source	<1 %
12	Submitted to Universitas Negeri Semarang Student Paper	<1 %
13	Y Hotmaida, Sutrisno. "Non-destructive determination of the main chemical components of red dragon fruit peel flour by using Near-Infrared Reflectance Spectroscopy (NIRS)", IOP Conference Series: Earth and Environmental Science, 2020 Publication	<1 %
14	Khalifa E. Sharquie. "Lactic Acid as a New Therapeutic Peeling Agent in Melasma",	<1 %

Dermatologic Surgery, 03/21/2006

Publication

15

Submitted to Ngee Ann Polytechnic

Student Paper

<1 %

16

www.ejournal.warmadewa.ac.id

Internet Source

<1 %

17

jppipa.unram.ac.id

Internet Source

<1 %

18

jurnal.utu.ac.id

Internet Source

<1 %

19

Dibyendu Kamilya, Mitila Debbarma, Prasenjit Pal, Biswanath Kheti, Sudipto Sarkar, Sukham Tushiba Singh. "Biofloc technology application in indoor culture of *Labeo rohita* (Hamilton, 1822) fingerlings: The effects on inorganic nitrogen control, growth and immunity", *Chemosphere*, 2017

Publication

<1 %

20

journal.poltekkes-mks.ac.id

Internet Source

<1 %

21

Edia Rahayuningsih, Felix Arie Setiawan, Ahmad Badawi Kasyfur Rahman, Tomimoto Siahaan, Himawan Tri Bayu Murti Petrus. "Microencapsulation of betacyanin from red dragon fruit (*Hylocereus polyrhizus*) peels using pectin by simple coacervation to

<1 %

enhance stability", Journal of Food Science and Technology, 2020

Publication

22

Gusti Ayu Mayani Kristina Dewi, Made Wirapartha, I K. A. Wiyana, N.P. M. Suartiningih. " Quality of eggs from fermented dragon fruit () flour fed Isa Brown ", IOP Conference Series: Materials Science and Engineering, 2020

Publication

<1 %

23

journal2.uad.ac.id

Internet Source

<1 %

24

Nina Meilisza, Muhammad Agus Suprayudi, Dedi Jusadi, Muhammad Zairin Jr., I Made Artika. "EFFECTS OF SYNTHETIC ASTAXANTHIN, CHLORELLA, AND SPIRULINA SUPPLEMENTATION IN DIETS ON GROWTH AND PIGMENTATION OF KURUMOI RAINBOWFISH, *Melanotaenia parva*", Indonesian Aquaculture Journal, 2021

Publication

<1 %

25

doc-pak.undip.ac.id

Internet Source

<1 %

26

fapet.ipb.ac.id

Internet Source

<1 %

27

ijeijournal.com

Internet Source

<1 %

28 repository.ubl.ac.id <1 %
Internet Source

29 www.e3s-conferences.org <1 %
Internet Source

30 Sri Wahdaningsih, Shoma Rizkifani, Eka Kartika Utari. "ANTI-AGING PEEL-OFF MASK OF DRAGON FRUIT PEEL EXTRACT (HYLOCEREUS POLYRHIZUS)", Jurnal Farmasi Sains dan Praktis, 2023 <1 %
Publication

31 www.jurnal.unsyiah.ac.id <1 %
Internet Source

Exclude quotes On

Exclude matches Off

Exclude bibliography On